

# “Possible Symptoms and Antifungal Treatment of Dermatitis Versicolor During a Population Found in Hot Status”

**Hailelassie Gebremeskel**

Department of Biotechnology, college of biological and chemical engineering, Mekelle University, Ethiopia.

**Issue Date:** March 04,2020 **Accepted Date:** August 23,2021 **Published Date:** September 03,2021

\***Corresponding Author:** 1Hailelassie gebremeskel, Department of Biotechnology, college of biological and chemical engineering, Mekelle University, Ethiopia. Tel: 924239947; E-mail: hailebiotek27@gmail.com, hailenebiland21@gmail.com

**Citation:** Gebremeskel H, Department of Biotechnology, college of biological and chemical engineering, Mekelle University, Ethiopia. J Pharm Microbiol. Vol:7.No.2

## Abstracts

Dermatitis versicolor is superficial fungal infection which is found as a commensal in several healthy adults. The foremost normally affected sites embody higher trunk, upper arms, neck and therefore the abdomen. Lesions confined to the acral elements like hands and feet have seldom been reported. Sun exposure, use of oils on the skin, naturally oily skin and sweating area unit all suspected to be triggers which will cause the spherical or oval yeast to convert to its hyphae sort, resulting in the rash and this infection isn't connected with poor hygiene. This text addresses many problems associated with characteristic the doable symptoms and creating the right identification and providing applicable treatment for this infection. Dermatitis versicolor is treated with antifungal medicines. These area units out there as shampoos, creams and tablets.

**Key words:** Dermatitis versicolor, Dermatitis, Malassezia species, Skin rash, Zymosis

## Introduction

Pityriasis versicolor could be a superficial zymosis of the skin caused by the yeast of the genus *Malassezia* and presents as hypo or hyper pigmented scaly macules.1 dermatitis versicolor (PV) conjointly known as tinea versicoloris a gentle however continual superficial zymosis of horny layer. This infection presents most ordinarily as pigmentation changes within the skin however can also be attended with cebure or pruritis.

The most usually affected sites embody higher trunk, upper arms, neck and also the abdomen. *Malassezia* could be a monophyletic genus of fungi found on the skin of seven billion humans and related to a spread of conditions, as well as dandruff, dermatitis (AE)/dermatitis, dermatitis versicolor, seborrheic eczema, and redness.

*Malassezia* species (formerly *Pityrosporum oval*) area unit completely the only eukaryotic member that forms a neighborhood of the resident skin small flora of human and different homothermic vertebrates with wide intra species diversity.

This intra-species diversity might be attributed to flora evolution because of ecology, host adaptation, and pathogenicity. These area unit expedient yeasts requiring predisposing environmental circumstances like temperature and humidness, patient immune standing, and genetic status. *Malassezia* spp. area unit related to completely different superficial pathologic process, as well as dermatitis versicolor (PV), seborrheic eczema (SD), redness still as healthcare facility blood infection in medical specialty care units.

Many species exist during this genus whereas the foremost common isolated kind from skin lesions is *Malassezia globosa*.

Many predisposing factors are projected for this illness like late immature and young adulthood age tropical and subtropics climate, temperate climate, use of oral contraceptives, poor hygiene and some different conditions.

## What is Malassezia Genus?

*Malassezia* genus is that the traditional mycoflora on human body covering surfaces. They colonize the regions containing oleaginous glands just like the head, neck and shoulders of humans. The genus of *Malassezia* has been divided into fourteen varied species supported the last categorization revision. *Malassezia* species area unit related to a large spectrum of clinical signs, like Tinea versicolor, redness and seborrheic eczema. It conjointly causes general infections in immune-compromised patients.

## Ecology and status of malassezia spices

*Malassezia* area unit super molecule dependent fungus yeasts that inhabit the skin and tissue layer sites of humans and different homothermic animals.

With the notable exception of *Malassezia pachydermatis*, all renowned *Malassezia* species need outwardly provided lipids for growth. It's been shown that this super molecule dependency is because of a defect within the synthesis of saturated fatty acid, that is the precursor of long-chain fatty acids.

This super molecule layer looks to safeguard M. furfur from phagocytosis, and down regulates the inflammatory reaction. Additionally, adhesion to host cells is also mediate by the property of the lipid-rich cell membrane.

The ability to metabolize lipids and to integrate the fatty acids into the flora cell membrane is crucial for growth and survival during host surroundings and contributes notably to the pathogenicity of *M. furfur*. Therefore, the enzymes necessary for these activities will be thought-about as virulence factors.

The unstable morphological variations, strict super molecule dependency result in issue in growing this plant, thereby limiting study on this plant.

Reclassification of this genus enclosed seven species particularly *M. furfur*, *M. pachydermatis*, *M. sympodialis*, *M. globosa*, *M. obtusa*, *M. restricta* and *M. slooffiae*.

*Malassezia* species were additionally found in wet subtropical climate and region that peopled by population with wide socio-cultural diversity as well as varied ethnic teams and migrant population of various elements at a time.

**Table 1:** Skin dermatosis caused or aggravated by infection by malassezia includes:

No	Type of Infection	Means of Occurrences
1		Most frequently because of the race <i>M. globosa</i> , <i>M. sympodialis</i> and <i>M. furfur</i>
2	Malassezia redness	Due to the yeast growing within the hair follicles wherever they manufacture inflammation
3	Seborrhoeic eczema	Dandruff, seborrheic and facial or scalp disease of the skin most frequently because of <i>M. restricta</i> and <i>M. globosa</i>
4	Neonatal cephalic pustulosis	A pustular eruption on young babies that resembles infantile disease of the skin

The plant life grows best in heat and wet conditions, explaining the upper prevalence of dermatosis versicolor in wet tropical climates.

There is a big increase in unwellness prevalence between childhood and adolescence, in all probability because of secretion changes that increase secretion production and permit for a lot of lipid-rich atmosphere during which the plant life will grow.

*Malassezia* species were originally thought to be specifically related to class hosts, however culture-independent studies discovered they'll occur in a very a lot of broader diversity of habitats, together with terrestrial and marine ecosystems like sea sediments, (Antarctic) soils, corals, sponges, nematodes, and cone snails.

Furthermore, *Malassezia* DNA was detected from soil nematodes in Central European forests, and it's been hypothesized that nematodes might function a vector for *Malassezia* species.

A population burst of *Malassezia* that proliferate extravagantly below favorable environmental conditions (e.g., increased heat, humidity). Reversing the annoying unwellness conditions will promptly resolve PV with none specific medical aid.

The unwellness is additionally a lot of common among adolescents and young adults WHO area unit physically active.

Pityriasis versicolor may be a harmless disease of the skin that causes scaly patches that typically itch over the trunk, neck, arms and legs. Anyone will grasp, however it's commonest in folks ages ten to thirty five. The rash ordinarily affects the trunk however is widespread, and also the face is especially affected in youngsters. The skin lesions area unit usually well or gently pruritic however is severely pruritic in bound wet conditions.

### Wherever this dermatosis versicolor is occurred

Dermatosis versicolor is additionally referred to as Tinea versicolor is common year-around within the tropics and climatic zone and is seen within the summer months in additional temperate climates. Sun exposure, use of oils on the skin, naturally oily skin and sweating area unit all suspected to be triggers that may cause the spherical or oval yeast to convert to its hyphae kind, leading to the rash. This rash will unfold between those that have skin contact.

The unstable morphological variations, strict macromolecule dependency cause problem in growing this plant life, thereby limiting study on this plant life. *Malassezia* species area unit tough to grow within the laboratory therefore scrapings is also reported as "culture negative". The yeast grows best if a macromolecule like vegetable oil is side to Littman agar matter.

### However the malassezia species cause infection on skin

The yeast lives in our pores. Below bound conditions, it will shift its kind from a spherical or oval yeast form to a string-like, branching form. These branching forms area unit named hyphae. The yeast will migrate below the skin and manufacture azelaic acid, a substance that may modification the quantity of pigment (color) in new skin cells. In its hyphae kind, the yeast causes a rash referred to as tinea versicolor, conjointly referred to as dermatosis versicolor.

Tinea versicolor happens a lot of typically in teenagers and young adults and seems on the rear, neck, higher chest, shoulders, armpits and higher arms. The general public WHO get this rash area unit in physiological state. However, tinea versicolor is a lot of doubtless to occur if you have got a suppressed system, for example, if you're taking a steroid medication like Deltas one for one more ill health. It's conjointly a lot of common in ladies WHO area unit taking contraception pills and in pregnant ladies.

### Symptoms of dermatosis versicolor

The roseola is created of scattered pink, tan, brown or white patches on the skin. All and sundry can typically have only 1 of the colors. These patches are typically flat, while not texture however might flake. They will be little spherical spots or areas

that begin smaller then get larger and mix with alternative areas. The rash could also be fretful, particularly once an individual is hot or sweating. Patches of skin could also be darker or lighter than your traditional skin color, or could also be red, brown or pink. They have an inclination to develop bit by bit and will link up to make larger patches over time.

The areas most frequently stricken by dermatosis versicolor embrace the rear, chest, upper arms, neck and tummy. This yeast is found on the skin of over ninetieth of adults, wherever it usually lives while not inflicting any issues. However dermatosis versicolor will develop if this yeast starts to multiply over usual. It isn't clear specifically why this happens in some folks and not in alternative.



Fig1: Pityriasis versicolor symptoms

### Interference of dermatosis versicolor

Because the rash happens additional promptly in hot, wet conditions, keep your skin as dry and funky as doable once you are during this climate. Tinea versicolor will be unfold through skin-to-skin contact or by coming back in grips with contaminated articles like towels, consumer goods or bedding.

### Many factors will increase your risk of developing dermatosis versicolor includes

- Living or staying in a very heat, dampish surroundings
- sweating to a fault (hyperhidrosis)
- creams, dressings or consumer goods that don't enable your skin to breathe
- being foodless, being a young person or in your early
- having a weakened system.<sup>43,44</sup>

Pityriasis versicolor isn't associated with poor hygiene. The condition cannot be unfold from person to person as a result of the majority have already got the *Malassezia* yeast on their skin.

### However dermatosis versicolor is treated

Pityriasis versicolor will be treated with antifungal medicines. These are out there as shampoos, creams and tablets.

### Antifungal shampoos

Antifungal shampoos (such as ketoconazole or atomic number 34 sulfide shampoo) are typically the primary treatment suggested for dermatosis versicolor.

In most cases, these shampoos got to be created into lather and left on the affected areas of skin for five to ten minutes

before being rinsed off. This typically has to be perennial each day for five to seven days. You will expertise some skin xerotes or irritation once mistreatment these shampoos, notably atomic number 34 sulfide. It should be useful to dilute the shampoo with water before applying it. Some folks additionally notice the odor of atomic number 34 sulfide shampoo unpleasant.

### Antifungal creams

If solely little areas of skin are affected, a medico might inflict associate antifungal cream. These creams typically got to be applied to the affected space of skin once or double daily for many weeks. Some folk's expertise a burning sensation once they use these antifungal creams, however this is often not common.

### Antifungal tablets

If an outsized space of skin is affected or alternative treatments haven't helped, you will be prescribed antifungal tablets. These typically got to be taken once daily for one to four weeks.<sup>46</sup> facet effects of those tablets are uncommon, though some folks expertise issues like rashes, feeling sick and tummy (abdominal) pain whereas taking them.

### Varieties of antifungal medicines

#### Antifungal medicines are out there as:

Topical antifungals – a cream, gel, ointment or spray you'll apply on to your skin, hair or nails Oral antifungals – a capsule, pill or liquid medication that you just swallow. Intravenous associated fungals – an injection into a vein in your arm, typically given in hospital. Intra epithelial duct antifungal pessaries – little, soft tablets you'll insert into the canal.

### Conclusions

*Malassezia* spices were additional found in wet climatic zone climate and region that colonized by population with wide socio-cultural diversity as well as varied ethnic teams and migrant population of various components at a time. several predisposing factors are planned for this illness like late teenaged and young adulthood age tropical and climatic zone climate, temperate climate, use of oral The flora grows best in heat and wet conditions, explaining the upper prevalence of dermatosis versicolor in wet tropical climates contraceptives, poor hygiene and many alternative conditions. Exposure to sun, using of oils and lotion on the skin, naturally oily skin and sweating are abilities to be triggers that may cause the spherical or oval yeast to convert to its hyphae type, leading to the rash. This rash will unfold between those that have skin contact. *Malassezia* species are tough to grow within the laboratory therefore scrapings could also be reportable as "culture negative". The yeast grows best if a macromolecule like vegetable oil is another to Littman agar medium. Dermatitis versicolor isn't largely transmitted from one individual as a result of the *malassezia* species are already gift on the skin of every individual. As a result of the rash happens additional pronto in

hot, wet conditions, keep your skin as dry and funky as doable after you area unit during this climate.

This dermatosis versicolor isn't connected with poor personal hygiene however is occurred in heat and wet environment; throughout overly sweating and dressing garments that aren't able to pass air in to the skin and with weakened system. This infection cannot be unfold from person to person as a result of most of the people have already got the *Malassezia* yeast on their skin.<sup>45</sup>

## Reference

- Arif T, Rather S (2015). cyst dermatosis versicolor-Rare variant of a typical medicine sickness, 6(1):105-106.
- Fitzpatrick T, Johnson RA, Wolff K, Suurmond D, editors. Color atlas and précis of clinical medicine. New York: Mc Graw Hill; 2001
- Saunders CW, Scheynius A, Heitman J (2012) *Malassezia* Fungi area unit specialized to live on Skin and regarding Dandruff, Eczema, and different Skin Diseases. PLoS Pathog 8(6): e1002701.
- Mishra et al. Exploration of anti-*Malassezia* potential of *Nyctanthes arbor-tristis* L. and their application to combat the infection caused by cheekbone s1 a totally unique matter. BMC Complementary and practice of drugs (2016) 16:114
- Velegraki A, Cafarchia C, Gaitanis G, Iatta R, Boekhout T. *Malassezia* infections in humans and animals: Pathophysiology, detection, and treatment. PLoS Pathog. 2015; 11(1):e1004523
- Gupta AK, Bluhm R, Summerbell R. dermatosis versicolor. J Eur Acad Dermatol Venereol. 2002; 16:19–33.
- Tarazooie B, Kordbacheh P, Zaini F, Zomorodian K, Saadat F, Zeraati H, et al. Study of the distribution of *Malassezia* species in patients with dermatosis versicolor and healthy people in capital of Iran, Iran. BMC Dermatol.2004; 4:5. Doi: 10.1186/1471-5945-4-5.
- Gaitanis G, Magiatis P, Hantschke M, Bassukas ID, Velegraki A. The *Malassezia* genus in skin and general diseases. Clin Microbiol Rev. 2012; 25:106–41.
- Tarazooie B, Kordbacheh P, Zaini F, Zomorodian K, Saadat F, Zeraati H, et al. Study of the distribution of *Malassezia* species in patients with dermatosis versicolor and healthy people in capital of Iran, Iran. BMC medicine 2004; 4:5.
- Salahi-Moghaddam A, Davoodian P, Jafari A, Nikoo MA. analysis of dermatosis versicolor in prisoners: A crosssectional study. Indian J Dermatol Venereol Leprol 2009; 75:379-82
- <https://www.medicalnewstoday.com/articles/315779#Treatment%20andprevention>
- <https://emedicine.medscape.com/article/1091575-overview>
- <https://www.cochranelibrary.com/es/cdsr/doi/10.1002/14651858.CD011208/full/es>
- Gemmer CM, DeAngelis YM, Theelen B, Boekhout T, Dawson TL. Fast, noninvasive methodology for molecular detection and differentiation of *Malassezia* yeast species on human skin and application of the strategy to dandruff biology. J Clin Microbiol. 2002; 40: 3350–3357.
- Porro, M. N., Passi, S., Caprill, F., Nazzaro, P. & Morpurgo, G. (1976). Growth necessities and lipide metabolism of *Pityrosporum orbiculare*. J Invest Dermatol sixty six, 178–182.
- Ashbee, H. R. & Evans, E. G. (2002). medicine of diseases regarding *Malassezia* species. Clin Microbiol Rev fifteen, 21–57
- Kesavan, S., Holland, K. T. & Ingham, E. (2000). the results of lipide extraction on the immunomodulatory activity of *Malassezia* species in vitro. Master of Education Mycol thirty eight, 239–247
- Mittag, H. (1995). Fine structural investigation of *Malassezia furfur*. II. The envelope of the yeast cells. Mycoses 38, 13–21
- Mancianti, F., Rum, A., Nardoni, S. & Corazza, M. (2001). living thing accelerator activity of *Malassezia* spp. isolates. Mycopathologia 149, 131–135.
- Sharma, A et al.2017, Characterization of *Malassezia* Species Isolated From dermatosis Versicolor Patients And Healthy Subjects of North-East India By PCR-RFLP And 26SrDNA Sequencing. Int J Recent Sci Res. 8(4), pp. 16624-16631
- Gupta AK, Yatika K, Summerbell RC. (2000): Molecular differentiation of seven *Malassezia* species. J Clin Microbiol, 38:1869-1875
- Cafarichia C, Gasser RB, Figueredo LA, Latrofa MS, Otranto D. (2011): Advances in identification of *Malassezia*. mole Cell Probes, 25:1-7
- Hellgren L, Vincent J. The incidence of tinea versicolor in central Scandinavian country. J Master of Education Microbiol 1983; 16: 501-2.
- Szepietowski JC, Baran E, Wild E. Tinea versicolor: a perspective study. Korean J Master of Education Mycol 2000;5: 108-12.
- semiotician metal, Jones MDM, Leonard G, Bass D. Marine fungi: their ecology and molecular diversity. Ann Rev Mar Sci. 2012; 4: 495–522.
- Amend A. From dandruff to sea vents: *Malassezia*-like fungi area unit ecologically hyper-diverse. PLoS Pathog. 2014; 10: e1004277.
- Renker C, Alpei J, Buscot F. Soil nematodes regarding the vertebrate moribific genus *Malassezia* (Basidiomycota: Ustilaginomycetes) in Central European forests. Biol Fertil Soils. 2003; 37: 70–72.
- Gupta AK, Bluhm R, Summerbell R. dermatosis versicolor. J Eur Acad Dermatol Venereol. 2002; 16: 19–33.

29. Kyriakis KP, Palamaras STI, Pagana G, Michailides C, Emmanuelides S. dermatosis versicolor prevalence by age and gender. *Mycoses* 2006; 49: 517-8.
30. Schwartz RA. Superficial plant infections. *Lancet* 2004; 364: 1173-82.
31. Gaitanis G, Magiatis P, Hantschke M, Bassukas ID, Velegriaki A. The *Malassezia* genus in skin and general diseases. *Clin Microbiol Rev* 2012; 25: 106-41.
32. [https://www.health.harvard.edu/a\\_to\\_z/tinea-versicolor-a-to-zor](https://www.health.harvard.edu/a_to_z/tinea-versicolor-a-to-zor) <https://www.aad.org>
33. Ashbee, H.R. (2006). Recent developments within the medicine and biology of *Malassezia* species. *FEMS medicine and Medical biology*, 47, 14-23.
34. Aspres, N., & Anderson, C. (2004). *Malassezia* yeasts within the pathological process of atopic dermatitis. *archipelago Journal of medicine*, 45, 199-207.
35. [https://www.health.harvard.edu/a\\_to\\_z/tinea-versicolor-a-to-z](https://www.health.harvard.edu/a_to_z/tinea-versicolor-a-to-z)
36. 14. Aspres, N., & Anderson, C. (2004). *Malassezia* yeasts within the pathological process of atopic dermatitis. *archipelago Journal of medicine*, 45, 199-207
37. [https://www.health.harvard.edu/a\\_to\\_z/tinea-versicolor-a-to-zor](https://www.health.harvard.edu/a_to_z/tinea-versicolor-a-to-zor) <https://www.aad.org>
38. Baker, B.S. (2006). The role of microorganisms in atopic dermatitis. *Clinical and Experimental medicine*, 144, 1-9.
39. [https://www.health.harvard.edu/a\\_to\\_z/tinea-versicolor-a-to-zor](https://www.health.harvard.edu/a_to_z/tinea-versicolor-a-to-zor) <https://www.aad.org>
40. <https://www.mayoclinic.org/diseases-conditions/tinea-versicolor/symptoms-causes/syc-20378385?page=0&citems=10>
41. Baroni, A., Paoletti, I., Ruocco, E., Agazzino, M., Tufano, M.A., & Donnarumma, G. (2004). doable role of *Malassezia* furfur in psoriasis: Modulation of TGF-beta one, integrin, and HSP70 expression in human keratinocytes and within the skin of psoriasis-affected patients. *Journal of animal tissue Pathology*, 31, 35-42
42. [https://www.health.harvard.edu/a\\_to\\_z/tinea-versicolor-a-to-z](https://www.health.harvard.edu/a_to_z/tinea-versicolor-a-to-z)
43. Bergman, J.N., & Eichenfield, J.N. (2002). infant disease of the skin and cephalic pustulosis. *Archives of medicine*, 138, 255- 257
44. *Malassezia dermatis* Sugita, Takashima, Nishikawa & Shinoda (2002)
45. Hu SW, Bigby M. dermatosis versicolor: a scientific review of interventions. *Arch Dermatol* 2010; 146: 1132-40.
46. *Malassezia yamatoensis* Sugita, Tajima, Takashima, Amaya, Saito, Tsuboi & Nishikawa (2004)
47. Gupta AK, Lane D, Paquet M. Systematic review of general treatments for tinea versicolor and evidence-based dosing programme recommendations. *J Cutan Master of Education Surg* 2014; 18:79-90.
48. Us Food and Drug Administration. Drug safety communication: office limits usage of Nizoral (ketoconazole) oral tablets due to doubtless fatal liver injury and risk of drug interactions and endocrine gland issues. 2013. [www.fda.gov/drugs/drugsafety/ucm362415.htm](http://www.fda.gov/drugs/drugsafety/ucm362415.htm)
49. <https://www.nhs.uk/conditions/pityriasis-versicolor/>
50. Bernier, V., Weill, F.X., Hirigoyen, V., Elleau, C., Feyler, A., Labreze, C., et al. (2002). Skin organization by *Malassezia* species in neonates: A prospective study and relationship with infant cephalic pustulosis. *Archives of medicine*, 138, 215-218.
51. Coutinho SD, Fedullo JD, Correa SH. Isolation of *Malassezia* spp. from wax of untamed felids. *Med Mycol.* 2006; 44(4): 383-7. Doi: 10.1080/13693780500411006